## HIV & AIDS and STI National Strategic Plan



## District-level data from priority districts (District Health Information System)

## Nevirapine uptake among new babies born to HIVpositive women

	Province	District	Nevirapine uptake rate among new babies born to HIV-positive women (%)				
			2003/04	2004/05	2005/06	2006/07	2007/08
1	Eastern Cape	Cacadu	117.1	87.1	103.5	22.1	99.5
2	Eastern Cape	Amathole	100.5	61.1	77.8	76.2	84.5
3	Eastern Cape	Chris Hani	79.6	87.8	85.6	13.2	93.6
4	Eastern Cape	Ukhahlamba	27.3	63.0	139.9	29.2	83.0
5	Eastern Cape	O.R. Tambo	193.1	140.2	90.2	46.6	95.0
6	Eastern Cape	Alfred Nzo	42.0	85.9	111.0	17.9	79.8
7	Free State	Thabo Mofutsanyane	46.9	61.3	97.3	65.7	104.9
8	Gauteng	Metsweding	-	-	66.7	19.4	100.0
9	KwaZulu-Natal	Amajuba	-	28.0	- 1	56.2	109.9
10	KwaZulu-Natal	Zululand	-	-		58.0	101.4
11	KwaZulu-Natal	Umkhanyakude	-	-		65.1	82.3
12	KwaZulu-Natal	iLembe	-	-	-	59.6	97.1
13	Limpopo	Mopani	95.0	98.0	96.6	67.4	99.7
14	Mpumalanga	Ehlanzeni	-	-	99.5	60.9	108.1
15	Northern Cape	Kgalagadi	100.0	333.3	98.6	65.7	101.4
16	North West	Bojanala Platinum	- 1	101.6	91.8	42.8	102.1
17	North West	Dr Ruth Segomotsi Mompati (Bophirima)	-	106.0	101.6	55.3	101.4
18	Western Cape	City of Cape Town	-	-	100.0	100.9	97.0

Source:

DHIS extracted

DHIS extracted

DHIS extracted

DHIS extracted DHIS

extracted

NOTE:

This indicator reflects the percentage of babies born to HIV-positive women, who receive Nevirapine suspension within 72 hours of birth. The numerator includes babies born at home as well as those born in a facility. For this reason the rates may be close to or over 100%

Downloaded from HST District Health Barometer www.hst.org.za

