

District-level data from priority districts
(District Health Information System)

Nevirapine uptake among HIV-positive pregnant women

	Province	District	Nevirapine uptake rate among HIV-positive pregnant women (%)				
			2003/04	2004/05	2005/06	2006/07	2007/08
1	Eastern Cape	Cacadu	94.6	69.4	76.2	85.3	80.0
2	Eastern Cape	Amathole	67.4	83.9	97.2	42.4	59.4
3	Eastern Cape	Chris Hani	23.8	43.0	57.4	49.0	67.7
4	Eastern Cape	Ukhahlamba	22.2	37.4	45.5	46.9	61.3
5	Eastern Cape	O.R. Tambo	37.1	43.2	40.1	33.2	50.9
6	Eastern Cape	Alfred Nzo	36.0	42.7	34.3	29.0	46.2
7	Free State	Thabo Mofutsanyane	5.4	24.2	73.3	63.4	25.4
8	Gauteng	Metsweding	2.1	0.0	72.6	71.6	59.7
9	KwaZulu-Natal	Amajuba	45.3	51.2	49.8	57.2	103.7
10	KwaZulu-Natal	Zululand	57.2	56.3	50.8	82.1	83.5
11	KwaZulu-Natal	Umkhanyakude	42.7	57.6	67.5	78.2	84.4
12	KwaZulu-Natal	iLembe	48.8	62.6	70.0	50.7	98.8
13	Limpopo	Mopani	36.7	35.5	50.7	55.7	62.5
14	Mpumalanga	Ehlanzeni	35.8	44.4	68.4	75.8	93.3
15	Northern Cape	Kgalagadi	25.0	9.6	29.3	64.5	100.4
16	North West	Bojanala Platinum	0.0	36.3	30.8	42.1	63.6
17	North West	Dr Ruth Segomotsi Mompati (Bophirima)	0.0	36.3	33.9	52.3	54.4
18	Western Cape	City of Cape Town	-	-	71.1	72.5	65.3

Source: DHIS extracted Oct 2008 DHIS extracted Oct 2008 DHIS extracted Oct 2008 DHIS extracted Oct 2008 DHIS extracted Oct 2008

NOTE:

This indicator reflects the proportion of HIV positive pregnant women who were dispensed with NVP to take home AND/OR those to whom NVP was administered at the facility during labour.

Downloaded from HST District Health Barometer

www.hst.org.za