SOUTH AFRICAN EARLY CHILDHOOD **REVIEW 2016**

CHILDREN BORN



6 300 000

4 000 000

KNOWLEDGE

INFORMATION & DATA SOLUTIONS

1MII OF THESE CHILDREN LIVE IN POVERTY IN SA EACH YEAR CHILDREN UNDER 6 IN SA (LESS THAN R923/MONTH) 12% of SA's children< 6 20 % of SA's children < 6 74% live in poverty 39% live in poverty LΡ 86% Rural 98% Urban 2% Rural 7% of SA's children < 6 71% live in poverty <u>8% of</u> SA's children< 6 ۰G MP 59% Rural NW 41% IIrha 64% live in poverty 62% Rura FS **KZN** NC 2% of SA's children< 6 21% of SA's children < 6 63% live in poverty 75% live in poverty 23% Rura 77% Urba 60% Rura 40% Ilrha EC 5% of SA's children < 6 WC 66% live in poverty 11% of SA's children < 6 83% Ilrhar 17 % Rural 42% live in poverty 14% of SA's children < 6 78% live in poverty 97% Urban 3% Rural 61% Rura

Living conditions for children under 6 in SA

LIVING IN **LIVING IN** LIVING WITH **URBAN AREAS** POVERTY **POOR SANITATION** Percentage of young children Percentage of young children Percentage of young children living living in urban areas has increased living in poverty has declined with poor sanitation has declined from 48% in 2002 to 57% in 2014 from 79% in 2003 to 63% in 2014 from 53% in 2002 to 26% in 2014 79% 63% 48% 53% 2002 2014 2003 2014 2002 2014

The **5** essential components



Maternal and child primary health (MCH) interventions

A child's health begins with the health of her mother. MCH services are critical for a safe pregnancy and birth, and to protect children Nutritional support

Good nutrition in pregnancy and early childhood is linked to better lifelong health and educational outcomes.

from disease during their most vulnerable developmental period.

WHAT'S NEEDED?

WHY IS THIS ESSENTIAL?

HOW IS SA DOING?

• A health facility within Adequate nutrition for pregnant women 30 minutes of every home Micronutrient supplementation • Early antenatal booking Deworming (before 20 weeks into pregnancy) Education on breastfeeding and child nutrition • HIV testing, prevention and treatment • Income support • Safe delivery in health facilities Full immunisation $\langle \cdot \rangle$ The percentage of women giving birth at (x) Anaemia in women of reproductive age is a health facilities has increased from 66% in major problem with severe consequences for 2002 to 86% in 2014. mother and child. Nearly a ¼ of SA women are $\langle \cdot \rangle$ The percentage of children completely anaemic. immunised by their 1st birthday has (X) Only 58% of infants aged 4 - 8 weeks are increased from 70% in 2002 to 90% in 2014. exclusively breastfed. Mother to child transmission of HIV is (x) 29% of poor children under 6 years in SA down to 2,6% nationally for infants aged 8 suffer from stunting. Stunting impacts both weeks. physical and cognitive development and is (X) Just half of all pregnant women have their associated with poor educational and health first antenatal visit before 20 weeks of outcomes later in life. pregnancy. All pregnant women must understand the Every clinic visit is an opportunity to reach importance of early antenatal care, and visit caregivers and children with nutritional a clinic within the first 3 months of their information and services. pregnancy. This is a gateway to timeous South Africa needs an anti-stunting campaign primary health services for them and their focused on achieving zero stunting within babies. the next ten years, which should include routine tracking of every child's growth and an immediate intervention when there are signs of growth faltering.

of the comprehensive package

3		5
Support for primary caregivers	Social services	Stimulation for early learning
Children need caregivers who are responsive and nurturing, from pregnancy, through babyhood, toddlerhood and beyond. Young children benefit from services targeting their mothers.	Social services and income support are critical to address child poverty and improve education, health and nutrition outcomes for children.	The foundations for lifelong learning are built in the first 6 years of life when the brain is most receptive to stimulation. Children who do not attend quality early learning programmes, start formal schooling at a disadvantage. The gap between them and their more advantaged peers widens over time.
In order to provide responsive and nurturing care, caregivers themselves need information and support, including parenting programmes and psychosocial services.	 Registration of birth within 30 days Immediate access to the child support grant (CSG) for eligible infants Early intervention for children who are at risk of neglect or abuse 	Early learning is supported in the home, in playgroups and early childhood development (ECD) centres or preschools.
 In 2014, 74% of mothers received follow up care 6 days after the birth of their child - a significant jump from just 5% in 2009. Antenatal and postnatal depression, and anxiety are huge problems affecting over 1/3rd of mothers in SA. SA has the highest rate of foetal alcohol syndrome in the world. On average, pregnant women are attending public antenatal facilities 4 times during pregnancy. 	 The number of children accessing the child support grant has surged from 22,000 in 1998 to 12 million in 2015. But, children are not getting the grant early enough, with just two out of three eligible infants under 1 year accessing the CSG. The child protection system is not equipped to deal with the high levels of physical and sexual abuse, and abandonment faced by young children in SA. 	 Access to early learning opportunities is unequal across income levels - 84% of children aged 3-5 years in the richest quintile have access, compared to just 57% in the poorest quintile. 1 million poor children aged 3-5 years do not have access to an early learning programme. For those children who do, programmes often don't meet minimum standards. Only 56% of Grade 3 learners achieve the 50% benchmarks for Maths and English. This is linked to poor access to quality early learning opportunities.
Antenatal and immunisation visits to health facilities provide up to 10 valuable opportunities to connect mothers to support during a critical period of their child's development. Maternal mental health and wellbeing need to be included as a routine part of these clinic interactions.	The uptake of antenatal care and high rates of delivery in health facilities are an opportunity for early birth registration and immediate enrolment on the child support grant. SA must strengthen the child protection system and ensure reliable data on the number of children needing social services.	The subsidy for early learning programmes needs to be increased to cover the full costs of a quality 20 hour per week programme, with sufficient support and monitoring to ensure quality.

Key indicators for early childhood development in South Africa



The South African Early Childhood Review 2016 is a joint publication between Ilifa Labantwana, the Children's Institute at the University of Cape Town and the Department of Planning, Monitoring and Evaluation in the Presidency.

The South African Early Childhood Review 2016 is an annual publication which presents information on the essential components of the comprehensive package of Early Childhood Development services. The data in this brief is drawn from a range of sources, which may be found in the full Review publication.

Ilifa Labantwana. ♥ PO Box 23893, Claremont 7735, 🔗 +27 (0)21 670 9848, 🖲 www.ilifalabantwana.co.za

This publication may be copied and shared. Please acknowledge the source.